

AD-A049 184

NORTHWESTERN UNIV EVANSTON ILL DEPT OF CHEMISTRY  
CHEMICAL AND STRUCTURAL RELATIONSHIPS AMONG THE OLIGOMERIC COMP--ETC(U)  
DEC 77 R D ERNST, T J MARKS

F/G 7/3  
N00014-77-C-0231

UNCLASSIFIED

TR-3

NL

| OF |  
AD  
A049 184



END

DATE  
FILMED

2-78

DDC



AD A 049184

AD No. \_\_\_\_\_  
DDC FILE COPY

OFFICE OF NAVAL RESEARCH

Contract N00014-77-C-0231

Task No. NR 053-640

TECHNICAL REPORT NO. 3

Chemical and Structural Relationships Among the Oligomeric Compounds  
 $MFe(CO)_4$  ( $M=Zn, Cd, Hg$ ),  $PbFe(CO)_4$ ,  $AgCo(CO)_4$ , and Their Lewis Base Adducts.

by

Richard A. D./Ernst  
Robin A. J./Marks

Prepared for Publication

in

Inorganic Chemistry

Northwestern University  
Department of Chemistry  
Evanston, Illinois 60201

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

Approved for Public Release; Distribution Unlimited

260 805

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 3	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Chemical and Structural Relationships Among the Oligomeric Compounds $MFe(CO)_4$ ( $M=Zn, Cd, Hg$ ), $PbFe(CO)_4$ , $AgCo(CO)_4$ , and Their Lewis Base Adducts		5. TYPE OF REPORT & PERIOD COVERED Interim, 1977
7. AUTHOR(s) R.D. Ernst and T.J. Marks		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Northwestern University, Evanston, IL 60201		8. CONTRACT OR GRANT NUMBER(s) N00014-77-C-0231
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Dept. of the Navy Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-053-640
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE December 29, 1977
		13. NUMBER OF PAGES 46
		15. SECURITY CLASS. (of this report) unclassified
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Fe - Zn bond      Ag - Co bond Fe - Cd Bond      Metal carbonyl Fe - Hg bond      Polymer Pb - Fe bond      Metal-metal bond		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  (see page 1)		

DDC  
JAN 27 1978  
RESOLVED  
F

DD FORM 1473  
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)



CHEMICAL AND STRUCTURAL RELATIONSHIPS AMONG  
THE OLIGOMERIC COMPOUNDS  $MFe(CO)_4$  ( $M = Zn, Cd, Hg$ ),  
 $PbFe(CO)_4$ ,  $AgCo(CO)_4$ , AND THEIR LEWIS BASE ADDUCTS.

by Richard D. Ernst<sup>1</sup> and Tobin J. Marks<sup>\*2</sup>

ABSTRACT

✓  
The synthesis and characterization of the following new metal-metal bonded complexes are reported:  $(pyridine)_3ZnFe(CO)_4$ ,  $(NH_3)(2,2'-bipyridyl)ZnFe(CO)_4$ ,  $(2,2':6,2''-terpyridyl)CdFe(CO)_4$ ,  $(2,2',2''-triaminotriethylamine)CdFe(CO)_4$ ,  $(2,2'-bipyridyl)ZnFe(CO)_4$ ,  $(4,4'-bipyridyl)CdFe(CO)_4$ , and  $(4-phenylpyridine)CdFe(CO)_4$ . Proper ligational strategy thus allows the isolation of several types of compounds which differ from the previously known  $B_2CdFe(CO)_4$  and  $(NH_3)_3ZnFe(CO)_4$  patterns. It is also possible to manipulate the degree of association of these species by control of the group IIb metal ion coordination sphere. Thus the first four of the new compounds listed are tentatively assigned monomeric structures, while the latter three are assigned oligomeric structures. The present studies also reveal that  $AgCo(CO)_4$  is isomorphous with  $[CdFe(CO)_4]_4$ , and that  $(pyridine)_2CdFe(CO)_4$  exists as a trimer in the solid state. The complex  $(2,2'-bipyridyl)AgCo(CO)_4$  is found to be partly associated in noncoordinating solvents. In the solid state,  $PbFe(CO)_4$  is associated.


↗

A

ACCESSION for	Wille Section	<input type="checkbox"/>	<input type="checkbox"/>
DATE	Burr Section	<input type="checkbox"/>	<input type="checkbox"/>
UNCLASSIFIED			
POST REPLY			
DO NOT WRITE IN THESE SPACES		SPECIAL	

✓

It has been known for some time that the metal-metal bonded oligomer  $[\text{CdFe}(\text{CO})_4]_n^3$  forms adducts,  $\text{B}_2\text{CdFe}(\text{CO})_4$ , with a wide variety of Lewis bases,  $\text{B}$ .<sup>3,4</sup> Until very recently there has been no structural information available for any of these materials. Diffraction studies have now shown that  $[\text{CdFe}(\text{CO})_4]_n$  is a tetramer with approximately  $D_{4h}$  symmetry<sup>5a</sup> and that  $(2,2'\text{-bpy})\text{CdFe}(\text{CO})_4$ , bpy = bipyridyl, is a trimer with approximately  $D_{3h}$  symmetry.<sup>5b</sup> The molecular geometries of these compounds are illustrated in Figure 1. The structures of  $\text{CdFe}(\text{CO})_4$  adducts with other Lewis bases remain largely undefined. The compound  $[\text{HgFe}(\text{CO})_4]_4$  is also known<sup>6,7</sup> and is isomorphous with  $[\text{CdFe}(\text{CO})_4]_4$ ,<sup>5a</sup> but no base adducts have been isolated. In comparison to the results for cadmium and mercury, the only reported zinc-containing derivative,  $\text{B}_3\text{ZnFe}(\text{CO})_4$  where  $\text{B} = \text{NH}_3$ ,<sup>7</sup> contains three molecules of base. Unlike the cadmium diammine,  $(\text{NH}_3)_2\text{CdFe}(\text{CO})_4$ , the zinc triammine does not readily lose ammonia on heating in vacuo.<sup>4</sup> A number of structural types are conceivable for the above Lewis base adducts, viz. A - C. In isoelectronic group Ib metal-transition metal carbonyl chemistry



$\text{M} = \text{Zn, Cd, Hg}$

an analogous range of possibilities exists for the compounds  $\text{AgCo}(\text{CO})_4$ ,  
 and  $\text{B}_n\text{AgCo}(\text{CO})_4$ .<sup>8</sup> Likewise, little is known about the related group  
 IVb derivative  $\text{PbFe}(\text{CO})_4$ .<sup>9</sup>

From the above discussion it can be seen that the relationships  
 between stable oligomeric structures as well as allowed structural  
 interconversions in group IIb and Ib metal-transition metal compounds  
 depend on ligands and metals in a way that has not been investigated in  
 depth. The details of, and reasons for, such relationships are of  
 importance to studies of metal-metal bonding in general and to the chal-  
 lenge of preparing infinite metal atoms arrays in particular. In this  
 article we report on the synthesis of several new types of group IIb  
 metal-transition metal complexes and discuss, on the basis  
 of physical measurements, possible chemical and structural relation-  
 ships in the  $\text{B}_n\text{MFe}(\text{CO})_4$  ( $\text{M} = \text{Zn}, \text{Cd}, \text{Hg}$ ),  $\text{AgCo}(\text{CO})_4$ ,  $\text{B}_n\text{AgCo}(\text{CO})_4$ , and  
 $\text{PbFe}(\text{CO})_4$  classes of molecules.



## EXPERIMENTAL

All operations involving metal carbonyls were carried out under an atmosphere of prepurified nitrogen in Schlenk apparatus or in a glove box. Solvents were thoroughly dried and deoxygenated in a manner appropriate to each, and were distilled under nitrogen immediately prior to use. Elemental analyses were performed by Miss H. Beck, Northwestern University Analytical Services Laboratory, Dornis and Kolbe Analytical Laboratories, or Schwarzkopf Analytical Laboratories. Osmometric molecular weight determinations were also carried out by Miss H. Beck using a Mecrolab Model 302 instrument.

Mercurytetracarbonyliron,  $\text{HgFe}(\text{CO})_4$ . This compound was prepared by the procedure of Brauer,<sup>10</sup> and was dried overnight under high vacuum. The infrared spectrum of a Nujol mull was identical with that reported previously.<sup>11</sup>

Thalliumtetracarbonylcobalt,  $\text{TlCo}(\text{CO})_4$ . This was prepared from thallium metal and  $\text{Co}_2(\text{CO})_8$  as described by Robinson *et al.*<sup>12a</sup> The infrared spectral parameters were identical to those previously reported,<sup>12b</sup> and are consistent with the essentially ionic structure found by x-ray structural determination<sup>12a</sup>.

Leadtetracarbonyliron,  $\text{PbFe}(\text{CO})_4$ . This very air-sensitive black compound was prepared by the method of Krumholz and Bril.<sup>9</sup>

Diamminecadmiumtetracarbonyliron,  $(\text{NH}_3)_2\text{CdFe}(\text{CO})_4$ . This compound was synthesized by the procedure of Feigl and Krumholz,<sup>3</sup> and was dried overnight under high vacuum. Infrared spectral parameters were identical with those previously reported.<sup>5a</sup>

Cadmiumtetracarbonyliron,  $\text{CdFe}(\text{CO})_4$  and bis(pyridine)cadmium tetracarbonyliron,  $(\text{py})_2\text{CdFe}(\text{CO})_4$ . These were prepared from  $(\text{NH}_3)_2\text{CdFe}(\text{CO})_4$  according to our previous procedures.<sup>5a</sup>

(2,2',2''-triaminotriethylamine)cadmiumtetracarbonyliron,  $(\text{tren})\text{CdFe}(\text{CO})_4$ . To a suspension of 1.40 grams (4.45 mmoles) of  $(\text{NH}_3)_2\text{CdFe}(\text{CO})_4$  in 40 ml diethylether under nitrogen was added an excess (1.0 ml) of 2,2',2''-triaminotriethylamine with stirring. After three hours, stirring was discontinued and the solid product was allowed to settle. The supernatant was removed, the product was washed three times with 25 ml portions of diethylether, and dried in vacuo overnight. The compound is only very slightly soluble in most organic solvents. Crystalline material could best be obtained by a controlled ligand replacement in which a solution of 2, 2', 2''- triaminotriethylamine is slowly reacted with (2,2'-bpy)<sup>5</sup> $\text{CdFe}(\text{CO})_4$ .

Anal. Calcd. for  $\text{C}_{10}\text{H}_{18}\text{CdFeN}_4\text{O}_4$ : C, 28.16; H, 4.25; N, 13.14.  
Found: C, 27.84; H, 4.31; N, 12.93.

Complete infrared data (Nujol mull): 3364m, 3310w, 3212m, 3126sh, 1948s, 1857s, 1815sh, 1798vs, br, 1583s, 1314m, 1268w, 1107w, 1094w, 1065s, 1037s, 1000m, 973m, 883m, 855w, 734w, 716w, 626s  $\text{cm}^{-1}$ .



Attempted preparation of bis(2,2'-bipyridyl)cadmiumtetracarbonyliron,  $(2,2'\text{-bpy})_2\text{CdFe}(\text{CO})_4$ . To 50 ml of deoxygenated aqueous ammonia was added 1.3 ml of  $\text{Fe}(\text{CO})_5$  (0.010 moles) under nitrogen and the resulting mixture was stirred overnight. To this solution of  $\text{HFe}(\text{CO})_4^-$  was added 50 ml of deoxygenated aqueous solution containing 2.67 grams (0.01 moles) of cadmium acetate dihydrate and 3.12 grams (0.02 moles) of 2-2' bipyridine at 0° C. This resulted in the precipitation of a yellow-orange solid which was filtered, washed with three 20 ml portions of deoxygenated water followed by three 20 ml portions of diethylether, and dried in vacuo. Elemental analysis (C,H,N), infrared spectroscopy, and x-ray powder diffraction identified the product as  $[(2,2'\text{-bpy})\text{CdFe}(\text{CO})_4]_3$ .<sup>5</sup>

2,2'-6',2''-terpyridylcadmiumtetracarbonyliron,  $(\text{tpy})\text{CdFe}(\text{CO})_4$ . A slight excess of 2,2':6',2''-terpyridine (0.20 grams, 0.86 mmoles) was dissolved in 10 ml acetone and added dropwise to 0.20 grams (0.71 mmoles) of  $\text{CdFe}(\text{CO})_4$  in 25 ml acetone under nitrogen with stirring. A reddish-orange precipitate immediately formed, and after two hours of stirring, the supernatant was removed. The solid product was then washed with three 25 ml portions of acetone, and dried in vacuo. The product so obtained smolders in air and is nearly insoluble in most organic solvents, except pyridine. Crystalline materials, significantly more air-resistant, may be obtained by slow diffusion of the reacting species through a medium glass frit.

Anal.<sup>13</sup> Calcd. for  $C_{19}H_{11}CdFeN_3O_4$ : C, 44.44; H, 2.16; N, 8.18, Cd, 21.89; Fe, 10.87. Found: C, 42.30; H, 2.48; N, 8.59, Cd, 22.32; Fe 11.20.

Complete infrared data (Nujol mull): 1985sh, 1973s, 1887s, 1872sh, 1850sh, 1593m, 1579w, 1573w, 1563w, 1433sh, 1419sh, 1306w, 1241w, 1186w, 1162w, 1155w, 998w, 754s, 645m, 622s  $cm^{-1}$ .

(4,4'-bipyridyl)cadmiumtetracarbonyliron, (4,4'-bpy)CdFe(CO)<sub>4</sub>.

This compound was prepared by stirring stoichiometric amounts of 4,4'-bipyridyl dihydrate (0.50 grams, 2.6 mmoles) and  $CdFe(CO)_4$  (0.73 grams, 2.6 mmoles) in 30 ml acetone under nitrogen for one day. The bright yellow precipitate which had formed immediately was then filtered from the supernatant, washed with several 10 ml portions of acetone, followed by several 20 ml portions of ether, and dried in vacuo.

Anal. Calcd. for  $C_{14}H_8CdFeN_2O_4$ : C, 38.52; H, 1.86; N, 6.42. Found: C, 37.83; H, 1.58; N, 6.03.

Complete infrared data (Nujol mull): 1977s, 1904s, 1892s, sh, 1604m, 1534w, 1491sh, 1414m, 1317w, 1219m, 1215m, 1062m, 1042w, 997m, 847w, 799m, 606s  $cm^{-1}$ .

(4-phenylpyridine)cadmiumtetracarbonyliron, (4-C<sub>6</sub>H<sub>5</sub>C<sub>5</sub>H<sub>4</sub>N)CdFe(CO)<sub>4</sub>.

To 0.33 grams (1.18 mmoles) of  $CdFe(CO)_4$  in 30 ml acetone at  $-78^\circ C$  under nitrogen was added 0.15 grams (0.97 mmoles) of 4-phenylpyridine with stirring. The solution was allowed to reach room temperature over a period of several hours, and was then stirred two more hours. The volume of the solution was next reduced by one-half, 50 ml pentane was added, and the solution cooled to  $-78^\circ C$ , producing a yellow mixture

of crystals and micro-crystalline powder. The supernatant was removed, the solid was washed twice with 20 ml portions of diethylether, and thoroughly dried in vacuo. The product was next taken up in a minimum volume of methylene chloride, and filtered from any residue (unreacted  $\text{CdFe}(\text{CO})_4$ ). An equal volume of pentane was added to the filtrate at  $-78^\circ\text{C}$ , producing a beautiful lemon-yellow micro-crystalline solid. After removal of solvent, the somewhat air-sensitive product was dried in vacuo for one hour.

Anal.: Calcd. for  $\text{C}_{15}\text{H}_{10}\text{CdFeNO}_4$ : C, 41.28; H, 2.31; N, 3.21.

Found: C, 40.87; H, 2.31; N, 3.33.

Complete infrared data (Nujol mull): 2000s, 1990sh, 1941sh, 1933sh, 1914s, 1907sh, 1891s, 1868sh, 1840m, 1806m, 1606m, 1546w, 1483sh, 1412m, 1219m, 1063m, 1040w, 1017w, 1003m, 912w, 827w, 756m, 748sh, 724w, 690w,  $605\text{s cm}^{-1}$ .

Tris(pyridine)zinctetracarbonyliron,  $(\text{py})_3\text{ZnFe}(\text{CO})_4$ . The compound  $(\text{NH}_3)_3\text{ZnFe}(\text{CO})_4$ , prepared as previously described,<sup>4,8</sup> was dissolved in an excess of deoxygenated pyridine under nitrogen. After evolution of  $\text{NH}_3$  was complete, a nitrogen flow was passed over the solution for one hour. Addition of ten volumes of pentane produced an oil, which solidified on cooling to  $-20^\circ\text{C}$ . The supernatant was removed, and the yellow-brown solid crushed to a powder. After washing with three 30 ml portions of diethylether, a bright yellow solid remained, which was dried in vacuo one hour.



Anal. Calcd. for  $C_{19}H_{15}FeN_3O_4Zn$ : C, 48.50; H, 3.21; N, 8.93.

Found: C, 48.50; H, 3.20; N, 8.33.

Complete infrared data (Nujol mull): 1974s, 1954s, 1879s, vbr, 1821s, vbr, 1796sh, 1603m, 1485sh, 1445sh, 1213m, 1153m, 1064m, 1036m, 1006m, 752m, 693s, 618s  $cm^{-1}$ .

(Ammine)(2,2'-bipyridyl)zinctetracarbonyliron,  $(NH_3)(2,2'-bpy)-ZnFe(CO)_4$ . To 50 ml aqueous ammonia was added 1.3 ml (0.01 mole) of  $Fe(CO)_5$  under nitrogen. After allowing this mixture to stir overnight (producing  $HFe(CO)_4^-$ ),<sup>4</sup> 50 ml of a deoxygenated aqueous solution containing 2.20 grams (0.01 mole) of zinc acetate dihydrate and 1.56 grams (0.01 mole) of 2,2'-bipyridyl was added at 0°C, causing a red-violet solid to appear. After one hour, the supernatant was removed and the solid was washed with three 30 ml portions of water, followed by three 30 ml portions of diethylether. The slightly air-sensitive product was then dried in vacuo one hour.

Anal. Calcd. for  $C_{14}H_{11}FeN_3O_4Zn$ : C, 41.37; H, 2.73; N, 10.34.

Found: C, 40.53; H, 2.63; N, 9.85.

Complete infrared data (Nujol mull): 3367m, 3346m, 3276w, 1975m, 1947s, 1914m, 1832s, vbr, 1788s, br, 1605m, 1596m, 1491sh, 1443sh, 1315m, 1204m, 1173w, 1157m, 1036w, 1020m, 1012sh, 834w, br, 760s, 726m, 643m, 625s  $cm^{-1}$ .

(2,2'-bipyridyl)zinctetracarbonyliron,  $(2,2'-bpy)ZnFe(CO)_4$ . By maintaining  $(NH_3)(2,2'-bpy)ZnFe(CO)_4$  at 75° in vacuo for three days, the

desired compound, of red-violet color, was obtained.

Anal. Calcd. for  $C_{14}H_8FeN_2O_4Zn$ : C, 43.18; H, 2.07; N, 7.19.

Found: C, 43.52; H, 2.28; N, 7.07.

Complete infrared data (Nujol mull): 2031w, 1975s, 1914s, 1881sh, 1866s, 1605m, 1597m, 1440sh, 1321m, 1258w, 1244w, 1172m, 1158m, 1060w, 1041m, 1015m, 1006sh, 760s, 727m, 715sh, 644m, 623s, 615sh  $cm^{-1}$ .

Silvertetracarbonylcobalt,  $AgCo(CO)_4$ . This was prepared by a variation of the method of Schubert,<sup>14</sup> wherein an ammoniacal aqueous  $AgNO_3$  solution is reacted with aqueous  $NaCo(CO)_4$ <sup>15</sup> under nitrogen. The yellow precipitate was then filtered, washed with pentane to remove  $Co_2(CO)_8$ , taken up in acetone, and filtered through a fine frit (to remove metallic silver). Addition of water to the filtrate caused precipitation of the compound. If necessary, the acetone/water recrystallization was repeated several times to complete the removal of silver (which, along with  $Co_2(CO)_8$  causes rapid catalytic decomposition of the compound).<sup>7, 15</sup> After removal of the supernatant, the solid compound was further washed with pentane, dried quickly (1 hr.) in vacuo, and stored in darkness at  $-20^\circ C$  to avoid decomposition.

Infrared data (Nujol mull): 2069s, 2034s, 2008s, 1993s, 1961s, br, 1937s  $cm^{-1}$ .

(2,2'-bipyridyl)silvertetracarbonylcobalt, (2,2'-bpy) $AgCo(CO)_4$ . This compound was prepared by a modification of the procedure of Hieber,



et al.<sup>8, 16</sup> The addition of 0.46 grams (2.95 mmole) of 2,2'-bipyridyl to a freshly prepared solution of  $\text{AgCo}(\text{CO})_4$  in ethanol (from the addition of 0.57 grams (2.95 mmole) of  $\text{NaCo}(\text{CO})_4$  in 30 ml ethanol to 0.50 grams (2.95 mmole) of  $\text{AgNO}_3$  in 30 ml ethanol) produced the desired compound. After stirring the mixture under nitrogen for 30 minutes, the solvent was removed in vacuo. The compound was then quickly extracted with a large volume of methylene chloride and the resulting solution was filtered, and reduced in volume under high vacuum. Addition of several volumes of pentane at  $-20^\circ \text{C}$  resulted in precipitation of the compound as a bright yellow solid.

Anal. Calcd. for  $\text{C}_{14}\text{H}_8\text{AgCoN}_2\text{O}_4$ : C, 38.65; H, 1.85; N, 6.44.

Found: C, 37.67; H, 1.80; N, 6.17.

Complete infrared data (Nujol mull): 2014s, 1967s, 1938s, 1910s, 1595sh, 1590m, 1566w, 1483sh, 1473sh, 1435sh, 1314m, 1303w, 1252w, 1240w, 1204w, 1169m, 1152m, 1097w, 1055w, 1036w, 1004m, 999sh, 747s, 730m, 635mw, 613w  $\text{cm}^{-1}$ .

Spectroscopic Studies. Infrared spectra were recorded with a Perkin Elmer 267 spectrophotometer. Mulls were prepared in a glove box with dry, degassed Nujol. Solutions were prepared in Schlenk apparatus and were transferred to matched NaCl cells as described previously.<sup>17</sup> All spectra were calibrated with polystyrene.

X-Ray Powder Diffraction Studies. X-ray powder diffraction patterns were recorded photographically on a Picker 6147 x-ray diffractometer

using filtered  $\text{CuK}\alpha$  radiation. Samples were sealed in 0.5 mm glass capillaries under nitrogen and rotated during exposure.

Unit Cell and Space Group Determination of bis(pyridine)cadmium-tetracarbonyliron,  $(\text{py})_2\text{CdFe}(\text{CO})_4$ . The compound was crystallized by very slow cooling of a hot saturated solution in pyridine/mesitylene (1:4), yielding large, well formed crystals. Elemental analysis of the single crystals proved them to be identical with the starting compound.<sup>18</sup>

A suitable single crystal was mounted under nitrogen in a glass capillary. Microscopic examination revealed no evidence of twinning. Oscillation, Weissenberg, and Precession photographs, using filtered  $\text{CuK}\alpha$  or  $\text{MoK}\alpha$  radiation revealed an orthorhombic unit cell,  $a = 12.42 \text{ \AA}$ ,  $b = 22.62 \text{ \AA}$ ,  $c = 17.04 \text{ \AA}$ ,  $V = 4788 \text{ \AA}^3$ . Systematic absences were observed for reflections of the type  $hk0$ ,  $h+k$  odd;  $h0l$ ,  $l$  odd;  $0kl$ ,  $k$  odd. These absences are uniquely consistent with space group #60,  $D_{2h}^{14}$ -Pbcn.<sup>19</sup> The observed density, 1.82 g/cc in a 1,4-dibromobutane/methylene bromide mixture agrees with the calculated value of 1.82g/cc for  $Z = 12$ . Since there are eight general positions in the unit cell, this means that, barring the unlikelihood of extreme disorder, the unit cell contains four trimeric units, which lie on two-fold axes (e.g., along  $0, y, \frac{1}{4}$ ).


## RESULTS

Group IIB Compounds


Previously reported group IIB metal-tetracarbonyliron complexes have been of the types  $MFe(CO)_4$  where  $M = Cd^3, Hg^{6,7}$ ;  $B_2MFe(CO)_4$  where  $M = Cd$  and  $B = NH_3^3$ , pyridine<sup>3</sup>,  $\gamma$ -picoline<sup>4</sup>,  $B_2 = 2,2'$ -bipyridyl<sup>4</sup>, 1,10-phenanthroline<sup>4</sup>, ethylenediamine<sup>4</sup>; as well as  $B_3MFe(CO)_4$  where  $M = Zn$  and  $B = NH_3^7$ . The scope of complexes which can be synthesized is expanded in the present work. By carefully controlling the stoichiometry, reaction conditions, and the nature of the ligands, we have obtained the following new types of compounds:  $BCdFe(CO)_4$ , from a low temperature reaction of  $B = 4-C_6H_5C_5H_4N$  with excess  $CdFe(CO)_4$ ;  $(4,4'$ -bpy) $CdFe(CO)_4$ , which may be polymeric by virtue of bridging nitrogenous ligands;  $B_3CdFe(CO)_4$ , by using the tridentate ligand 2,2': 6', 2''-terpyridyl (tpy); possibly  $B_4CdFe(CO)_4$  by using the potentially tetradentate ligand 2,2',2''-triaminotriethylamine (tren);<sup>20</sup> and finally,  $B_2ZnFe(CO)_4$  by careful removal of  $NH_3$  from  $(NH_3)(2,2'$ -bpy) $ZnFe(CO)_4$  in vacuo. Attempts to stabilize the  $B_4CdFe(CO)_4$  species  $(2,2'$ -bpy) $_2CdFe(CO)_4$  by coordination of the base on cadmium prior to reaction with  $HFe(CO)_4^-$  lead simply to loss of the excess base and isolation of  $[(2,2'$ -bpy) $CdFe(CO)_4]_3$ . Attempts to prepare base adducts of  $HgFe(CO)_4$  using mono- or bidentate nitrogen donor ligands also failed, as determined by x-ray powder diffraction, infrared spectroscopy, and elemental analysis. Synthetic procedures and analytical data are compiled in the Experimental Section.



From our previous work in group IIb<sup>5,21</sup> and group IVb<sup>22,23</sup> metal-transition metal chemistry as well from established electron density - CO force constant relationships in metal carbonyl vibrational spectroscopy<sup>5,24</sup>, it is possible to draw a number of qualitative structural inferences about the new Fe(CO)<sub>4</sub> derivatives from band positions, splittings and intensities in the infrared spectra. In particular, progressive coordination of Lewis base molecules to the main group metal is expected to weaken or destroy metal-metal bonding<sup>5,23,25</sup>. Furthermore, the accompanying base stabilization of ionic resonance hybrids such as D is predicted to



result in a lowering of C-O stretching frequencies due to increased population of C-O  $\pi^*$  orbitals and decreased C-O bond order. Whether a particular B<sub>n</sub>MFe(CO)<sub>4</sub> species is or is not oligomeric can be investigated by comparing spectra in the solid state<sup>24f</sup> or in noncoordinating solvents with those obtained in coordinating solvents (e.g. pyridine). Base-induced metal-metal bond fracture by the coordinating solvent will result in a shift of the  $\nu$ C-O absorptions to lower energy. Upon full cleavage to bimetallic units, the band pattern generally simplifies and is characteristic of the C<sub>3v</sub> local symmetry (2A<sub>1</sub>, 1E modes) monomer, F<sup>5</sup>.




Such structures have been identified for isoelectronic  $R_2(B)M'Fe(CO)_4$ <sup>23, 25a</sup> and  $X_3M'Fe(CO)_4$ <sup>- 26</sup> derivatives ( $M'$  = group IVb metal),  $(py)_3CdFe(CO)_4$  (shown to be monomeric in solution)<sup>5</sup>,  $HFe(CO)_4$ <sup>- 27</sup>, and  $RFe(CO)_4$ <sup>-</sup> ( $R = n-C_3H_7$ )<sup>28</sup>. Cleavage of all metal-metal bonds to yield an ionic structure, A, would displace the C-O vibrational frequencies to the region of  $Na_2Fe(CO)_4$ , i. e.  $1720\text{ cm}^{-1}$  in  $[(cryptate)Na]_2Fe(CO)_4$ ,<sup>28a</sup>  $1730\text{ cm}^{-1}$  in DMF,<sup>29</sup> and  $1761\text{ cm}^{-1}$  as a Nujol mull.<sup>27</sup> As will be seen, the above vibrational spectroscopic information can be supplemented by x-ray powder diffraction studies in the solid and molecular weight measurements in solution. Detailed metrical data, however, will only come from complete single crystal diffraction studies.

Solid state infrared spectra in the C-O stretching region of representative group IIb metal-transition metal derivatives are shown in Figure 2. Numerical data are summarized in Table I along with solution molecular weights. It can be seen that increasing the number as well as the basicity of donor molecules coordinated to the group IIb metal generally leads to



a lowering of the carbonyl stretching frequencies. Also, zinc complexes have lower energy C-O stretching transitions than cadmium analogues. These trends reflect increasing polarity of metal-metal bonding and negative charge build-up on the  $\text{Fe}(\text{CO})_4$  fragment. The lowest C-O stretching frequencies are observed for the  $\text{B}_3\text{MFe}(\text{CO})_4$  complexes. The energies of the carbonyl stretching absorptions in the latter compounds shift very little on dissolution in coordinating solvents. Furthermore, solution spectra (Figure 3) are in good agreement with the approximately  $\text{C}_{3v}$  structure F.<sup>5,24</sup> These observations viewed in light of the molecular weight data and the saturation of the  $\text{M}^{+2}$  coordination sphere strongly suggest monomeric structures<sup>30</sup>. The complex  $(\text{tren})\text{CdFe}(\text{CO})_4$  possesses the lowest energy  $\nu_{\text{C-O}}$  absorptions of any of the molecules examined. Though the energies and the molecular symmetry as revealed by the band multiplicity are not compatible with a purely ionic structure, it appears that tetradentate  $\text{Cd}^{+2}$  coordination by the tren ligand may be placing additional negative charge on the tetracarbonyliron moiety. In none of the compounds studied was the characteristic C-O band position of an isocarbonyl species<sup>31</sup>, G, observed. A strong transition in



the infrared spectrum of  $(\text{tren})\text{CdFe}(\text{CO})_4$  at  $1583\text{ cm}^{-1}$  can be assigned to an  $\text{NH}_2$  deformation<sup>32</sup>; in the free ligand this band occurs at  $1592\text{ cm}^{-1}$ .

The  $\text{B}_2\text{MFe}(\text{CO})_4$  molecules all appear to be associated to some degree. We have already shown  $(2, 2'\text{-bpy})\text{CdFe}(\text{CO})_4$  to be trimeric in the solid state and in noncoordinating solvents<sup>5</sup>. The space group and unit cell parameters of  $(\text{py})_2\text{CdFe}(\text{CO})_4$  also suggest a trimeric structure in the crystal (see Experimental Section for details). On the basis of infrared spectral and molecular weight studies, this compound is known to exist in pyridine solution as monomeric  $(\text{py})_3\text{CdFe}(\text{CO})_4$ <sup>5a</sup>. X-ray powder diffraction studies revealed that  $(2, 2'\text{-bpy})\text{ZnFe}(\text{CO})_4$  and  $(2, 2'\text{-bpy})\text{CdFe}(\text{CO})_4$  are not isomorphous. The exact structure of the Zn compound remains undetermined, however the similarity of the  $\nu_{\text{C-O}}$  multiplicities and energies to those of the Cd analogue strongly suggest an associated structure. This compound is insufficiently soluble for solution molecular weight determination. The  $\text{BCdFe}(\text{CO})_4$  compound  $(4\text{-C}_6\text{H}_5\text{C}_5\text{H}_4\text{N})\text{CdFe}(\text{CO})_4$  is found to be dimeric in benzene (Table I). Though a similar structure could be adopted by  $(4, 4'\text{-bpy})\text{CdFe}(\text{CO})_4$ , the very low solubility of this complex along with the lower C-O stretching frequencies suggests an oligomeric bpy cross-linked structure with both nitrogen atoms involved in  $\text{Cd}^{+2}$  coordination (to different cadmium ions).

### Silver and Lead Complexes

A number of structural similarities exist between isoelectronic pairs of group IIb and group IVb metal-transition metal complexes. For


example, the  $B_3MFe(CO)_4$  and  $R_2(B)M'Fe(CO)_4$ <sup>23, 25a</sup> ( $M' = Si, Ge, Sn, Pb$ ) compounds are largely monomeric, while  $B_2CdFe(CO)_4$  and  $R_2M'Fe(CO)_4$ <sup>33</sup> compounds appear, in the absence of overwhelming steric factors, to be associated. These similarities motivated the examination of relationships among similar compounds involving group IVb (divalent), group IIIb, and group Ib metal ions. For example, the structures of  $AgCo(CO)_4$ <sup>14</sup> and  $PbFe(CO)_4$ <sup>9</sup> are unknown, while those of their respective isoelectronic analogs,  $[CdFe(CO)_4]_4$ <sup>5a</sup> and  $TlCo(CO)_4$ <sup>12</sup> (largely ionic structure)<sup>12a</sup> are known. The solid state infrared spectra of  $PbFe(CO)_4$  and  $TlCo(CO)_4$  are compared in Figure 2, and are seen to be rather dissimilar. The spectrum of  $PbFe(CO)_4$  resembles that of an oligomeric structure, i. e. the bands are shifted very much to high frequency, coming near those of  $CdFe(CO)_4$  and  $Fe(CO)_4Cl_2$  (Table I), whereas the spectrum of the more ionic  $TlCo(CO)_4$  is reasonably close to that of  $NaCo(CO)_4$  (Table I, Figure 2). In agreement with these observations, x-ray powder patterns reveal that these compounds are not isomorphous. Efforts to cleave  $PbFe(CO)_4$  with a variety of Lewis bases were unsuccessful.

In contrast to the Pb, Tl results, the infrared spectra of  $AgCo(CO)_4$  and  $[CdFe(CO)_4]_4$  are qualitatively rather similar (Figure 2), although the C-O stretching transitions of  $AgCo(CO)_4$  occur at somewhat higher energy, in accord with the lower formal negative charge on the  $Co(CO)_4^-$  fragment as compared to the  $Fe(CO)_4^{2-}$  fragment<sup>24</sup>. This result suggests that the



compounds may be composed of similar oligomeric units. Indeed, x-ray powder powder diffraction data (Table II) show them to be isomorphous. Thus, in  $\text{AgCo}(\text{CO})_4$ , the metal atoms are also arrayed in an approximately planar, centrosymmetric eight-membered ring. Unlike  $[\text{CdFe}(\text{CO})_4]_4$ ,  $[\text{AgCo}(\text{CO})_4]_4$  is particularly prone to thermal decomposition, yielding Ag and  $\text{Co}_2(\text{CO})_8$ . The decomposition is catalyzed so greatly by even traces of Ag that various attempts to crystallize the compound by a reversible homolytic metal-metal bond cleavage cycle have so far been fruitless.

The compounds  $(2,2'\text{-bpy})\text{AgCo}(\text{CO})_4$  and  $[(2,2'\text{-bpy})\text{CdFe}(\text{CO})_4]_3$  were also investigated for a possible structural relationship. The Nujol mull and solution infrared spectra of these compounds (Table I and Figure 3) are not particularly similar, and the solid compounds, each recrystallized from  $\text{CH}_2\text{Cl}_2$ /hexane, are not isomorphous by x-ray powder diffraction. The infrared spectrum of  $(2,2'\text{-bpy})\text{AgCo}(\text{CO})_4$  in  $\text{CH}_2\text{Cl}_2$  (Figure 3) is essentially that expected for a monomeric  $C_{3v}$  structure such as H.<sup>24, 34</sup> Furthermore, the compound is found to be



H

only partially associated in methylene chloride solution by osmometry. Relatively small shifts of the C-O stretching transitions are observed in pyridine

solution (Table I, Figure 3) indicating a similar structure with possible weak coordination of an additional molecule of base. The related compound (o-triars)AgCo(CO)<sub>4</sub> (o-triars=bis(o-dimethylarsinophenyl)methylarsine) is known to be monomeric in solution and exhibits a similar infrared spectrum<sup>35</sup> (Table I).


## DISCUSSION

The results of these studies have shown that  $B_n M Fe(CO)_4$  complexes (for M = Zn or Cd) other than those normally favored by typical monodentate ligands in the solid state can be isolated using an appropriate ligational strategy. Thus, not only have solution-state structures (e.g., monomeric  $B_3 Cd Fe(CO)_4$ )<sup>5</sup> been stabilized in the solid state, but structures previously unknown in either phase have been obtained (e.g.,  $B_2 Zn Fe(CO)_4$  and  $B Cd Fe(CO)_4$ ). The solid state structures of these and isoelectronic species thus far observed fall into three categories of aggregation: ionic (A), monomeric (B), and oligomeric (C). Through this series the iron coordination number ranges from four to six. The ionic extreme has already been established in [(cryptate)Na]<sub>2</sub>Fe(CO)<sub>4</sub><sup>36</sup> as well as in isoelectronic TlCo(CO)<sub>4</sub>,<sup>12</sup> while (B)R<sub>2</sub>M'Fe(CO)<sub>4</sub>,<sup>23, 25a</sup>  $B_3 Cd Fe(CO)_4$ ,<sup>5</sup> and (o-triars)AgCo(CO)<sub>4</sub><sup>35</sup> are known to be monomeric. Similarly, oligomeric structures have been demonstrated for [CdFe(CO)<sub>4</sub>]<sub>4</sub>,<sup>5</sup>  $B_2 Cd Fe(CO)_4$ ,<sup>5</sup> and R<sub>2</sub>M'Fe(CO)<sub>4</sub> species<sup>33</sup> with normal R groups. The present studies



have shown that  $B_2ZnFe(CO)_4$  ( $B_2 = 2, 2'$ -bpy),  $BCdFe(CO)_4$  ( $B = 4-C_6H_5C_5H_4N$ ), and  $AgCo(CO)_4$  are most likely oligomeric, while  $B_3ZnFe(CO)_4$  ( $B = py, (NH_3)(2, 2'$ -bpy)),  $B_nAgCo(CO)_4$  ( $B_n = (2, 2'$ -bpy)), and  $B_3CdFe(CO)_4$  ( $B_3 = tpy, tren$ ) are most likely monomeric, or only very weakly associated.


Two key factors in determining which structure is favored for a given complex seem to be the formal positive charge on, and the electronegativity of, the main group metal. For example, in comparing the isoelectronic  $B_2AgCo(CO)_4$  and  $B_2CdFe(CO)_4$  complexes, the oligomeric nature of the latter species plausibly derives from the greater difficulty in separating ions of formal charge  $\pm 2$  as compared to  $\pm 1$  in the former complexes. Within group IIb, the most common compounds of zinc are monomeric with three base molecules ( $B_3ZnFe(CO)_4$ ) and those of cadmium are oligomeric in the solid state with two base molecules ( $B_2CdFe(CO)_4$ ), while  $HgFe(CO)_4$  does not form base adducts. This trend reflects the propensity of the harder and more acidic main group metals to be coordinated by the nitrogen donor ligands, which are harder than the  $Fe(CO)_4$  moiety. Alternatively, it could be said that zinc, being the most electropositive group IIb metal, is best capable of stabilizing the positive charge resulting from resonance hybrid I. Species of the type  $B_3CdFe(CO)_4$  are



generally observed only when very large excesses of B are present (e. g. in

pyridine solutions). However, even though zinc displays a strong preference for bonding to the harder nitrogen donors in monomeric complexes, we have prepared the  $(2,2'\text{-bpy})\text{ZnFe}(\text{CO})_4$  oligomer from the  $(\text{NH}_3)(2,2'\text{-bpy})\text{ZnFe}(\text{CO})_4$  monomer by taking advantage of the greater volatility of  $\text{NH}_3$ . Our inability to isolate base adducts of  $\text{HgFe}(\text{CO})_4$  parallels previous work on  $\text{M}[\text{Co}(\text{CO})_4]_2$  compounds ( $\text{M} = \text{Zn}, \text{Cd}, \text{Hg}$ ).<sup>37</sup> The marked tendency for mercury to retain two-coordination has often been attributed to a high 6s - 6p promotional energy.<sup>38</sup>

After considering the factors which control stoichiometry and the extent of association of the complexes, it is next appropriate to discuss the extent of oligomerization in some of the specific cases. From the unit cell and space group determination of  $(\text{py})_2\text{CdFe}(\text{CO})_4$  (see Experimental Section), we have found that this complex is *trimeric in the solid state*, as in the case of the related 2,2'-bipyridyl complex<sup>5b</sup>. In contrast, the isoelectronic  $\text{R}_2\text{SnFe}(\text{CO})_4$  compounds are dimeric<sup>33</sup> (Structure J). The trimeric structures of  $(2,2'\text{-bpy})\text{CdFe}(\text{CO})_4$  and  $(\text{py})_2\text{CdFe}(\text{CO})_4$



appear to reflect the preferred valence angles at the group IIb metal and at iron. The Fe- Cd-Fe angles in the bipyridyl complex are in the range  $138.81(15)^\circ$  -  $148.40(15)^\circ$ ,<sup>5b</sup> while the Fe-Sn-Fe angles in  $[(CH_3)_2SnFe(CO)_4]_2$ <sup>33b</sup> are  $103.7(4)^\circ$  and  $102.6(3)^\circ$ , and in  $[(\eta^5-C_5H_5)_2SnFe(CO)_4]_2$ <sup>33a</sup> are  $102.04(3)^\circ$ . The larger angles in the Cd compound seem to reflect a preference for greater cadmium s character in the more covalent Cd-Fe bonds (as opposed to the weaker Cd-N coordinate bonds), while in the group IVb system the Sn-C and Sn-Fe bonds are more equal in covalency and the hybridization is expected to be more near tetrahedral. It should be noted, in support of relatively constant cadmium hybridization, that in the series  $B_nCd[Mn(CO)_5]_2$ , B = terpyridyl,<sup>39a</sup> diglyme,<sup>39b</sup> 2,2'-bipyridyl,<sup>39c</sup> and 1,10-phenanthroline<sup>39c</sup>, the observed Mn-Cd-Mn angles are in the relatively narrow range of  $128-136^\circ$ . The larger Fe-Cd-Fe angles in  $B_2CdFe(CO)_4$  species clearly favor a ring size larger than 4 members. In addition to factors involving the main group metal ion, it was noted in the discussion of the  $[(2,2'-bpy)CdFe(CO)_4]_3$  structure<sup>5b</sup> that the tetracarbonyliron coordination geometry was appreciably distorted from octahedral toward bicapped tetrahedral (the two cadmium atoms located at the capping vertices), with Cd-Fe-Cd angles ranging from  $94.9(1)^\circ$  to  $102.2(2)^\circ$  and  $C_{axial}-Fe-C_{axial}$  angles of  $135.8(15)^\circ$ - $141.8(15)^\circ$ . This distortion was attributed to the large amount of electron density being released by the cadmium atom onto the  $Fe(CO)_4$  fragment. Such a distortion also destabilizes a 4-membered ring both because the Cd-Fe-Cd angles would be too large and because close non-bonded contacts ( $\sim 2\text{\AA}$ )



are predicted between axial carbonyl ligands on opposite iron atoms.

In contrast, Sn-Fe-Sn angles in the two aforementioned group IVb compounds were reported to be  $76.3(4)^\circ$ ,  $77.4(3)^\circ$ ; <sup>33b</sup> and  $77.96(2)^\circ$ , <sup>33a</sup> respectively. This contraction from the octahedral  $90^\circ$  <sup>40</sup> is clearly more compatible with a 4-membered ring. Not surprisingly, infrared spectra in the C-O stretching region reveal less electron density on the iron in the group IVb than in the group IIb compounds <sup>5</sup>. That the new group IIb compound  $[(4-C_6H_5C_5H_4N)CdFe(CO)_4]_2$  is apparently dimeric in solution suggests that, to maintain a planar ring structure, the tetracarbonyliron valence angles must decrease markedly in response to the removal of a base molecule from cadmium, while the Fe-Cd-Fe angles must decrease to less than  $120^\circ$ . A solid state structure determination for this compound would clearly be of interest.

#### Isoelectronic Relationships


In our studies on the two isoelectronic pairs,  $TlCo(CO)_4$  and  $PbFe(CO)_4$ , and  $AgCo(CO)_4$  and  $CdFe(CO)_4$ , the conclusions drawn on the basis of infrared spectroscopy and x-ray powder diffraction parallel other known relationships. Thus, the first pair are as different in their chemical and physical properties as they are in their spectroscopic ones. While  $TlCo(CO)_4$  is yellow, sublimable, and soluble in aromatic solvents, <sup>12</sup>  $PbFe(CO)_4$  is black, non-sublimable, insoluble in aromatic solvents, and only very slightly soluble in coordinating solvents. <sup>9</sup> Both are very air-sensitive, however. In this case, the chemical and physical differences

are probably related to the fact that while Pb(II) tends toward high coordination numbers involving a stereochemically active (hence, nonbonding) lone pair,<sup>41</sup> in Tl(I) the lone pair (at or near room temperature, at least) is essentially spherical, and hence metal-to-ligand sigma antibonding which, along with the lower formal charge on thallium, tends to reduce the tendency to high coordination numbers.<sup>42</sup>


The structural similarities of the second isoelectronic pair,  $\text{AgCo(CO)}_4$  and  $\text{CdFe(CO)}_4$  (both planar tetramers) fall in line with other chemical similarities. Both compounds are yellow, nonvolatile, soluble in coordinating solvents (e. g., acetone) from which they can be precipitated by the addition of water (cf., reversible homolytic cleavage cycle),<sup>5,23a</sup> and form bipyridyl adducts. The chemical similarities in this case seem to stem from the ability of the two isoelectronic main group metals, Ag(I) and Cd(II), to adopt either linear, two-coordinate  $sp$  complexes, or higher (especially three or four) coordination numbers.<sup>43</sup> However, the formal difference in charge does manifest itself in the 2,2'-bipyridyl adducts, where the cadmium complex is oligomeric while the silver complex is only partially associated. It can also be noted that while many cis-octahedral  $\text{Fe(CO)}_4\text{X}_2$  complexes are known,<sup>40</sup>  $[\text{AgCo(CO)}_4]_4$  appears to be the first such cobalt complex.

While  $\text{AgCo(CO)}_4$  and  $\text{CdFe(CO)}_4$  are isoelectronic and isostructural, some interesting differences do exist in their available resonance hybrids.



Those of  $\text{CdFe}(\text{CO})_4$  and its base adducts involve (neglecting ligands):



while the corresponding resonance structures for  $\text{AgCo}(\text{CO})_4$  and its base adducts are



For the silver-cobalt system, a much more convenient resonance formalism can be adopted, utilizing donation of lone pairs of electrons from the transition metal carbonyl to the main group metal. This is known in the related compounds  $(\text{C}_5\text{H}_5)\text{Co}(\text{CO})_2 \cdot \text{HgCl}_2$ ,<sup>44a</sup>  $(\text{C}_5\text{H}_5)\text{Co}(\text{olefin})_2 \cdot \text{HgCl}_2$ ,<sup>44a</sup>  $(\text{C}_5\text{H}_5)\text{Rh}(\text{olefin})_2 \cdot \text{HgCl}_2$ ,<sup>44a</sup>  $\text{Fe}(\text{CO})_5 \cdot \text{HgCl}_2$ ,<sup>44b</sup>  $[(\text{C}_6\text{H}_5)(\text{CH}_3)_2\text{P}]_2\text{Fe}(\text{CO})_3 \cdot \text{HgCl}_2$ <sup>44c</sup> as well as others.<sup>44d-h</sup> Thus,  $\text{H}:\overset{\text{H}}{\text{Fe}}(\text{CO})_4$ , in forming the dianion, clearly has at least two electron pairs available, as does the isoelectronic  $\text{Co}(\text{CO})_4^-$ . Resonance structures (K) and (L) (neglecting ligands) can then be invoked in which the extent of lone pair donation is variable.



### Concluding Remarks

In our synthetic, spectroscopic, and crystallographic studies on  $\text{B}_n\text{MFe}(\text{CO})_4$  and  $\text{B}_n\text{M}'\text{Co}(\text{CO})_4$  type compounds, we have found that the



main group metal with attendant ligands plays a dominant role in determining the type of structure adopted and the resulting chemical and physical properties. Besides determining its own coordination sphere and tendency to form base adducts, the main group metal is primarily responsible for the extent of cation/anion association in these compounds, which encompasses ionic, monomeric, oligomeric, and possibly (but not presently) isocarbonyl structures. By suitable adaptation of the main group metal and the ligands chosen, we have shown that otherwise unfavored structures for a particular metal can be stabilized, thus potentially allowing for more detailed analysis of formerly unstable or solution-state structures. Such manipulations could also result in preferential stabilization of infinite metal chains, and will be the subject of future investigations.

#### ACKNOWLEDGMENTS

We thank the Northwestern Materials Research Center (NSF DMR 72-0319A06) the Paint Research Institute, and the Office of Naval Research (N00014-77-C-0231) for generous support of this work. We thank Prof. J. A. Ibers for the use of x-ray equipment. We thank W. R. Grace and Company for generous gifts of dessicants.

## REFERENCES

1. NSF Predoctoral Fellow, 1974-1977.
2. a. Fellow of the Alfred P. Sloan Foundation.  
b. Camille and Henry Dreyfus Teacher-Scholar.
3. F. Feigl and P. Krumholz, Z. Anorg. Allg. Chem., 215, 242 (1933).
4. A. T. T. Hsieh, M. J. Mays, and R. H. Platt, J. Chem. Soc. A., 3296 (1971).
5. a. R. D. Ernst, T. J. Marks, and J. A. Ibers, J. Am. Chem. Soc., 99, 2090 (1977).  
b. R. D. Ernst, T. J. Marks, and J. A. Ibers, J. Am. Chem. Soc., 99, 2098 (1977).
6. (a) H. Hock and H. Stuhlman, Chem. Ber., 61, 2097 (1928);  
(b) Ibid., 62, 431 (1929).
7. W. Hieber and E. Fack, Z. Anorg. Allg. Chem., 236, 83 (1938).
8. W. Hieber and H. Schulten, Z. Anorg. Allg. Chem., 232, 17 (1937).
9. P. Krumholz and S. Bril, Proc. Chem. Soc., 116 (1960).
10. G. Brauer, "Handbook of Preparative Inorganic Chemistry", 2nd ed., Academic Press, New York, N. Y., 1963, Vol. II, p. 1755.
11. T. Takano and Y. Sasaki, Bull. Chem. Soc. Japan, 44, 431 (1971).
12. a. D. P. Schussler, W. R. Robinson, and W. F. Edgell, Inorg. Chem., 13, 153 (1974).  
b. J. M. Burlitch and T. W. Theyson, J. Chem. Soc. Dalton Trans., 828 (1974).
13. This compound is apparently fairly refractory and combustion analyses invariably yielded low carbon analyses, the best being 43.28%.

14. (a) M. P. Schubert, J. Am. Chem. Soc., 55, 4563 (1933);  
 (b) J. J. Eisch and R. B. King, Eds., "Organometallic Syntheses", Vol. I, Academic Press, New York, N. Y., 1965, p. 102.
15. W. F. Edgell and J. Lyford, IV, Inorg. Chem., 9, 1932 (1970).
16. W. Hieber, E. O. Fischer, and E. Böckly, Z. Anorg. Allg. Chem., 269, 308 (1952).
17. T. J. Marks, J. Chem. Educ., 48, 167 (1971).
18. Anal Calcd for  $C_{14}H_{10}CdFeN_2O_4$ : C, 38.34; H, 2.30; N, 6.39. Found: C, 38.57; H, 2.34; N, 6.09.
19. N. F. M. Henry and K. Lonsdale, Eds., "International Tables for X-ray Crystallography," Vol. I, Kynoch Press, Birmingham, England, 1969.
20. F. A. Cotton and G. Wilkinson, "Advanced Inorganic Chemistry", 3rd ed., Interscience, New York, N. Y., 1972, p. 625.
21. For general reviews of Group IIb metal-transition chemistry see:
  - a. M. J. Taylor, "Metal-to-Metal Bonded States of the Main Group Elements," Academic Press, N. Y., 1975, Chapt. 2.
  - b. A. T. T. Hsieh and M. J. Mays, MTP Int. Rev. Sci., Inorg. Chem., Ser. One, 6, 43 (1972).
  - c. B. P. Birynkov and Yu. T. Struchkov, Russ. Chem. Rev., 39, 789 (1970).
22. For general reviews of Group IVb metal-transition metal chemistry see:
  - a. Reference 21a, chapt. 4.
  - b. F. Höfler, Topics Curr. Chem. 50, 129 (1974).
  - c. E. H. Brooks and R. J. Cross, Organometal. Chem. Rev., A6, 227 (1970).
  - d. J. F. Young, Advan. Inorg. Chem. Radiochem., 11, 92 (1968).



23. a. T.J. Marks and A.R. Newman, J. Amer. Chem. Soc., 95, 769 (1973).
- b. G.W. Grynkewich, B.Y.K. Ho, T.J. Marks, D.L. Tamaja, and J.J. Zuckerman, Inorg. Chem., 12, 2522 (1973).
- c. For structural work on analogous  $R_2(B)SnCr(CO)_5$  systems see: F.A. Cotton and M.D. Brice, J. Amer. Chem. Soc., 95, 4529 (1973); T.J. Marks, J. Amer. Chem. Soc., 93, 7090 (1971).
24. a. P.S. Braterman, Struct. Bond., 26, 1 (1976).
- b. P.S. Braterman, "Metal Carbonyl Spectra", Academic Press, London, 1975.
- c. L.M. Haines and M.H.B. Stiddard, Adv. Inorg. Chem. Radiochem., 12, 53 (1969).
- d. G. Bor, Inorg. Chim. Acta, 1, 81 (1967).
- e. A. Reckziegel and M. Bigorgne, J. Organometal. Chem., 3, 341 (1965).
- f. Structural deductions derived from solid state spectral patterns are obviously complicated by site symmetry, correlation field, and other environmental effects<sup>24g,h</sup>. However the weighted average of the band positions can still serve as a qualitative index of the electron density on the transition metal, and features such as high ionic character can be clearly recognized.
- g. S.F.A. Kettle, Pure Appl. Chem., 27, 113 (1971).
- h. J.G. Bullitt and F.A. Cotton, Inorg. Chim. Acta., 5, 637 (1971).
25. a. A.B. Cornwell, P.G. Harrison, and J.A. Richards, J. Organometal. Chem., 108, 47 (1976).
- b. A. Davison and J.E. Ellis, J. Organometal. Chem., 36, 113 (1972).
- c. J.M. Burlitch, J. Amer. Chem. Soc., 91, 4562 (1969).
26. a. Th. Kruck and W. Molls. Z. Anorg. Allg. Chem., 420, 159 (1976).
- b. Th. Kruck, E. Job, and U. Klose, Angew. Chem. Int. Ed. Engl., 7, 374 (1968).
- c. Th. Kruck and B. Herber, Angew. Chem. Int. Ed. Engl., 8, 679 (1969).

27. K. Farmery, M. Kilner, R. Greatrex, and N. N. Greenwood, J. Chem. Soc., A, 2339 (1969).
28. a. J.P. Collman, R.G. Finke, J.N. Cawse, and J.I. Brauman, J. Amer. Chem. Soc. 99, 2515 (1977).  
 b. G. Hüttner and W. Gartzke, Chem. Ber., 108, 1373 (1975).
29. W.F. Edgell, M.T. Yang, B.J. Bulkin, R. Bayer, and N. Koizumi, J. Amer. Chem. Soc., 87, 3080 (1965).
30. a) The infrared spectrum of  $(py)_3ZnFe(CO)_4$  in  $CH_2Cl_2$  shows that a mixture of species is present, including  $(py)_3ZnFe(CO)_4$ . The other component shows C-O stretching modes at higher frequencies, and is thus possibly a  $(py)_2ZnFe(CO)_4$  complex arising from pyridine dissociation, i.e.,  $(py)_3ZnFe(CO)_4 \rightleftharpoons (py)_2ZnFe(CO)_4 + py$ .  
 b. It is recognized that the coplanarity of the three nitrogen atom donors in 2,2':6',2''-terpyridyl means that the  $Cd^{+2}$  ion coordinative saturation is not as extensive as in the  $B_3$  complexes with monodentate bases. Thus a possibility for oligomerization does exist,<sup>39a</sup> but such a situation is not unambiguously indicated in the infrared spectra of  $(tpy)CdFe(CO)_4$ .
31. a. D.F. Shriver, Acc. Chem. Res., 3, 231 (1970).  
 b. G.J. Gainsford, R.R. Schrieke, and J.D. Smith, J. Chem. Soc., Chem. Commun., 650 (1972).  
 c. A.E. Crease and P. Legzdins, J. Chem. Soc., Dalton Trans., 1501 (1973).  
 d. S.W. Ulmer, P.M. Skarstad, J.M. Burlitch, and R.E. Hughes, J. Am. Chem. Soc., 95, 4469 (1973).
32. K. Nakamoto, "Infrared Spectra of Inorganic and Coordination Compounds," 2nd ed., Wiley-Interscience, New York, N.Y., 1970, pp. 150-158.
33. In all cases examined to date, these complexes are dimeric for sterically normal R groups:  
 a. P.G. Harrison, T.J. King, and J.A. Richards, J. Chem. Soc. Dalton Trans., 2097 (1975).  
 b. C.J. Gilmore and P. Woodward, J. Chem. Soc., Dalton Trans., 1387 (1972).  
 c. J.C. Zimmer and M. Huber, C.R. Acad. Sci., Ser. C, 267, 1685 (1968).  
 d. G.W. Grynkewich and T.J. Marks, Inorg. Chem., 15, 1307 (1976), and references therein.
34. a. G.C. vandenBerg, A. Oskam, and K. Breize, J. Organometal. Chem., 69, 169 (1974).

34. b. G.C. van den Berg and A. Oskam, J. Organometal. Chem., 78, 357 (1974).
35. A.S. Kasenally, R.S. Nyholm, and M.H.B. Stiddard, J. Chem. Soc., 5343 (1965).
36. R.G. Teller, R.G. Finke, J.P. Collman, H.B. Chin, and R. Bau, J. Amer. Chem. Soc., 99, 1104 (1977).
37. a. W. Hieber and R. Breu, Chem. Ber., 90, 1259 (1957).  
b. J.M. Burlitch and A. Ferrari, Inorg. Chem., 9, 563 (1970).
38. Reference 20a, p. 504.
39. a. W. Clegg and P.J. Wheatley, J. Chem. Soc., Dalton Trans., 90 (1973).  
b. W. Clegg and P.J. Wheatley, J. Chem. Soc., Dalton Trans., 424 (1974).  
c. W. Clegg and P.J. Wheatley, J. Chem. Soc., Dalton Trans., 511 (1974).
40. a. The stereochemical nonrigidity observed in  $(R_3M)_2Fe(CO)_4$  compounds<sup>41b-e</sup> suggests considerable deformability of the  $M_2Fe(CO)_4$  coordination sphere along certain reaction coordinates.  
b. R.K. Pomeroy, L. Vancea, H.P. Calhoun, and W.A.G. Graham, Inorg. Chem., 16, 1508 (1977).  
c. L. Vancea, M.J. Bennett, C.E. Jones, R.A. Smith, and W.A.G. Graham, Inorg. Chem., 16, 897 (1977).  
d. L. Vancea, R.K. Pomeroy, and W.A.G. Graham, J. Amer. Chem. Soc., 98, 1407 (1976).  
e. T.J. Marks and G.W. Grynkewich, J. Organometal. Chem., 91, C9(1975).
41. P.G. Harrison, Coord. Chem. Revs., 20, 1 (1976).
42. A.G. Lee, Coord. Chem. Revs., 8, 1 (1972).



43. Reference 20, p. 505, 1046.
44. a. I.N. Nowell and D.R. Russell, Chem. Commun., 817(1967).  
b. H. Hock and H. Stuhlman, Chem. Ber., 61B, 2097(1928).  
c. Reference 21b, footnote 23.  
d. D.J. Cook, J.L. Dawes, and R.D.W. Kemmitt, J. Chem. Soc., A, 1548 (1967).  
e. D.M. Adams, D.J. Cook, and R.D.W. Kemmitt, J. Chem. Soc., 106 (1968).  
f. K. Edgar, B.F.G. Johnson, J. Lewis, and S.B. Wild, J. Chem. Soc., 2851 (1968).  
g. J.L. Dawes and R.D.W. Kemmitt, J. Chem. Soc., 1072 (1968).  
h. J.L. Dawes and R.D.W. Kemmitt, J. Chem. Soc., 2093 (1968).

TABLE I.

Comparison of Infrared-Active C-O Stretching Frequencies in Related Metal Carbonyl Complexes<sup>a</sup>

Compound	Phase	$\nu_{\text{CO}}, \text{cm}^{-1}$	Molecular Weight <sup>b</sup>	Solvent
$[\text{CdFe}(\text{CO})_4]_4$	Nujol mull	2032s, 2012m, 1972ms, 1961ms, 1946m, 1920s, br 1867s, br, 1840w, sh		
$\text{PbFe}(\text{CO})_4$	Nujol mull	2049s, 2006m, 1990w, sh 1978w, sh, 1964ms		
$[(4-\text{C}_6\text{H}_5\text{C}_5\text{H}_4)\text{CdFe}(\text{CO})_4]_2$	Nujol mull	2000s, 1990sh, 1941sh, 1933sh, 1914s, 1907sh, 1891s, 1868sh, 1840m, 1806m		
$[(4-\text{C}_6\text{H}_5\text{C}_5\text{H}_4\text{N})\text{CdFe}(\text{CO})_4]_2$	$\text{CH}_2\text{Cl}_2$	1992m, 1927s, br	819(873)	benzene
$[(2,2'\text{-bpy})\text{CdFe}(\text{CO})_4]_3^c$	benzene	1982vs, 1922s, 1894ms, 1879sh	1344(1308)	benzene
$(\text{py})_3\text{CdFe}(\text{CO})_4^c$	pyridine	1976s, 1908vs, 1886sh	320(280) <sup>c</sup>	pyridine
$(\text{tpy})\text{CdFe}(\text{CO})_4$	Nujol mull	1985sh, 1973s, 1887s, 1872sh, 1850sh		
$(\text{tpy})\text{CdFe}(\text{CO})_4$	pyridine	1977s, 1907vs, 1888sh		
$(\text{tren})\text{CdFe}(\text{CO})_4$	Nujol mull	1948s, 1857s, 1815sh 1798vs, br		

TABLE I continued

Compound	Phase	$\nu_{\text{CO}}, \text{cm}^{-1}$	Molecular Weight <sup>b</sup>	Solvent
(4,4'-bpy) <sub>2</sub> CdFe(CO) <sub>4</sub>	Nujol mull	1977s, 1904s, 1892s, sh		
(2,2'-bpy) <sub>2</sub> ZnFe(CO) <sub>4</sub>	Nujol mull	1975s, 1914s, 1881sh, 1866s		
(NH <sub>3</sub> )(2,2'-bipy)ZnFe(CO) <sub>4</sub>	Nujol mull	1975m, 1947s, 1914m, 1832s, 1788s		
(NH <sub>3</sub> ) <sub>3</sub> ZnFe(CO) <sub>4</sub>	Nujol mull	1994m, 1943s		
(py) <sub>5</sub> ZnFe(CO) <sub>4</sub>	Nujol mull	1974s, 1954s, 1879s, br, 1821s, 1796sh		
(py) <sub>3</sub> ZnFe(CO) <sub>4</sub>	pyridine	1954ms, 1857ms, 1823s	487(470)	pyridine
[Et <sub>4</sub> N]HFe(CO) <sub>4</sub> <sup>d</sup>	Nujol mull	2008mw, 1910sh, 1848s, br		
Na <sub>2</sub> Fe(CO) <sub>4</sub> <sup>24a</sup>	Nujol mull	1761s		
Na <sub>2</sub> Fe(CO) <sub>4</sub> <sup>24b</sup>	DMF	1730s		
[(cryptateNa) <sub>2</sub> Fe(CO) <sub>4</sub> ] <sup>e</sup>	KBr	1720s		
Fe(CO) <sub>4</sub> Cl <sub>2</sub> <sup>f</sup>	C <sub>2</sub> Cl <sub>4</sub>	2164w, 2124s, 2108m, 2084s		



TABLE I continued

Compound	Phase	$\nu_{\text{CO}}$ , $\text{cm}^{-1}$	Molecular Weight <sup>b</sup>	Solvent
$\text{AgCo}(\text{CO})_4$	Nujol mull	2069s, 2034s, 2008s 1993s, 1961s, br, 1937s		
$\text{TiCo}(\text{CO})_4$	Nujol mull	2020mw, 1931vs, 1900m, sh	376(375) <sup>g</sup>	benzene <sup>g</sup>
$(2,2'\text{-bpy})\text{AgCo}(\text{CO})_4$	Nujol mull	2014s, 1967s, 1938s, 1910s		
$(2,2'\text{-bpy})\text{AgCo}(\text{CO})_4$	$\text{CH}_2\text{Cl}_2$	2023ms, 1954sh, 1940s, br	603(435)	
$(2,2'\text{-bpy})\text{AgCo}(\text{CO})_4$	pyridine	2011ms, 1930sh, 1910s, br		
$(\text{o-triars})\text{AgCo}(\text{CO})_4$ <sup>h</sup>	$\text{CHCl}_3$	2037s, 1953sh, 1923s	778(730) <sup>h</sup>	benzene
$\text{NaCo}(\text{CO})_4$ <sup>i</sup>	DMF	1892s		
$\text{Co}(\text{CO})_4\text{Cl}$ <sup>j</sup>	acetone	2120m, 2068m, 2046s, br		

<sup>a</sup>m = medium; s = strong; sh = shoulder; br = broad; w = weak; v = very.

<sup>b</sup>Value in parenthesis is that calculated for the formula unit shown in the left-hand column.

<sup>c</sup>Reference 5. <sup>d</sup>Reference 27. <sup>e</sup>Reference 28a.

<sup>f</sup>K. Noack, Helv. Chim. Acta., **45**, 1847 (1962).

<sup>g</sup>S.R. Pedersen and W.R. Robinson, Inorg. Chem., **14**, 2360 (1975).

<sup>h</sup>Reference 35. <sup>i</sup>W.F. Edgell, M.T. Yang, and N. Koizumi, J. Am. Chem. Soc., **87**, 2563(1965).

<sup>j</sup>M. Pankowsky and M. Bigorgne, Compt. Rend., **C264**, 1382(1967).

TABLE II

Most Prominent X-Ray Powder Diffraction Lines for


 $[\text{CdFe}(\text{CO})_4]_4$  and  $\text{AgCo}(\text{CO})_4$ , mm. ( $1\text{mm}/^\circ 4\theta$ )

$[\text{CdFe}(\text{CO})_4]_4$	$\text{AgCo}(\text{CO})_4$
18.4s	18.0s
20.4w	19.9m
29.2s	28.0s
36.6s	34.5s, 36.1mw
43.6mw	43.2m
62.1s	60.8s
--	67.1m
72.5s	72.0s


Figure 1

Perspective views of  $[(2,2'\text{-bpy})\text{CdFe}(\text{CO})_4]_3$  (from reference 5b) and  $[\text{CdFe}(\text{CO})_4]_4$ , (from reference 5a).





$[\text{CdFe}(\text{CO})_4]_4$



$[(\text{bpy})\text{CdFe}(\text{CO})_4]_3$

Figure 2

Nujol Mull Infrared Spectra of Various Metal Carbonyl Complexes.

- A.  $\text{AgCo(CO)}_4$
- B.  $\text{TlCo(CO)}_4$
- C.  $\text{PbFe(CO)}_4$
- D.  $(2, 2'\text{-bpy})\text{ZnFe(CO)}_4$
- E.  $(\text{py})_3\text{ZnFe(CO)}_4$
- F.  $\text{CdFe(CO)}_4$
- G.  $(4\text{-C}_6\text{H}_5\text{C}_5\text{H}_4\text{N})\text{CdFe(CO)}_4$
- H.  $(4, 4'\text{-bpy})\text{CdFe(CO)}_4$
- I.  $(\text{tpy})\text{CdFe(CO)}_4$
- J.  $(\text{tren})\text{CdFe(CO)}_4$



















Figure 3

Solution infrared spectra for various monomeric metal-metal bonded carbonyls.

- A.  $(\text{tpy})\text{CdFe}(\text{CO})_4$  in pyridine
- B.  $(\text{py})_3\text{ZnFe}(\text{CO})_4$  in pyridine
- C.  $(2,2'\text{-bpy})\text{AgCo}(\text{CO})_4$  in  $\text{CH}_2\text{Cl}_2$
- D.  $(2,2'\text{-bpy})\text{AgCo}(\text{CO})_4$  in pyridine



# TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
Office of Naval Research Arlington, Virginia 22217 Attn: Code 102IP 1	6	U.S. Army Research Office P.O. Box 12211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. Jerry Smith	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Weapons Center China Lake, California 93555 Attn: Head, Chemistry Division	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Civil Engineering Laboratory Port Hueneme, California Attn: Mr. W. S. Haynes	1
ONR Branch Office 760 Market Street, Rm. 447 San Francisco, California 94102 Attn: Dr. P. A. Miller	1	Professor O. Heinz Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
ONR Branch Office 495 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Director, Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Office of Naval Research Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1
The Asst. Secretary of the Navy (R&D) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1		
Commander, Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1		



# TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. R. M. Grimes University of Virginia Department of Chemistry Charlottesville, Virginia 22901	1	Dr. W. Hatfield University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514	1
Dr. M. Tsutsui Texas A&M University Department of Chemistry College Station, Texas 77843	1	Dr. D. Seyferth Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1
Dr. C. Quicksall Georgetown University Department of Chemistry 37th & O Streets Washington, D.C. 20007	1	Dr. M. H. Chisholm Princeton University Department of Chemistry Princeton, New Jersey 08540	1
Dr. M. F. Hawthorne University of California Department of Chemistry Los Angeles, California 90024	1	Dr. B. Foxman Brandeis University Department of Chemistry Waltham, Massachusetts 02154	1
Dr. D. B. Brown University of Vermont Department of Chemistry Burlington, Vermont 05401	1	<del>Dr. T. Marks Northwestern University Department of Chemistry Evanston, Illinois 60201</del>	<del>1</del>
Dr. W. B. Fox Naval Research Laboratory Chemistry Division Code 6130 Washington, D.C. 20375	1	Dr. G. Geoffrey Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1
Dr. J. Adcock University of Tennessee Department of Chemistry Knoxville, Tennessee 37916	1	Dr. J. Zuckerman University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1
Dr. A. Cowley University of Texas Department of Chemistry Austin, Texas 78712	1		